

Isoniazid

eye-soe-NYE-a-zid

Brand Name: Nydrazid

Drug Class: Opportunistic Infection and Other Drugs

Isoniazid, also known as INH, belongs to the class of medicines called antimycobacterials. These medicines prevent or treat infections caused by a certain type of bacterium, including the bacterium that causes tuberculosis (TB).

HIV/AIDS-Related Uses

Isoniazid is approved by the FDA for the prevention and treatment of TB in people with HIV/AIDS, either by itself or in combination with other antimycobacterials.

Dosage Form/Administration

Isoniazid comes in oral (syrup or tablet) and injectable forms. The oral forms should be taken either 1 hour before or 2 hours after a meal, but may be taken with food if necessary to avoid gastrointestinal distress. Aluminum hydroxide-containing antacids decrease absorption of isoniazid, so isoniazid must be taken at least 1 hour before the antacid.

Isoniazid is available combined with other anti-TB medicines into one pill: with rifampin in a capsule called Rifamate, or with rifampin and pyrazinamide in a tablet called Rifater. Another drug, pyridoxine, is usually given together with isoniazid to help prevent some of the nervous system side effects caused by isoniazid.

Contraindications

Individuals should tell a doctor about any medical problems before taking this medicine.

Possible Side Effects

Along with its desired effects, isoniazid can cause some unwanted effects, including hepatitis (dark urine, yellow eyes or skin); hepatitis prodromal symptoms (loss of appetite, nausea or vomiting, unusual tiredness or weakness); peripheral neuritis (clumsiness or unsteadiness; numbness; tingling; burning; or pain in hands and feet); seizures; mood or mental changes; skin rash or sores; fever; unusual bleeding or bruising; and blurred vision or loss of vision, with or without eye pain. Individuals should tell a doctor if they have any of these symptoms.



Other side effects may not be serious and may lessen or disappear with continued use of this medicine. Less serious side effects of this medicine include stomach pain, diarrhea, nausea, and vomiting. Individuals should tell a doctor if these side effects continue or are bothersome.

Drug and Food Interactions

A doctor should be notified of any other medications being taken, including prescription, nonprescription (over-the-counter), or herbal medications.

Clinical Trials

For information on clinical trials that involve Isoniazid, visit the ClinicalTrials.gov web site at <http://www.clinicaltrials.gov>. In the Search box, enter: Isoniazid AND HIV Infections.

Manufacturer Information

Nydrazid
Sandoz Inc
506 Carnegie Center Drive
Suite 400
Princeton, NJ 08540
(609) 627-8500

Isoniazid
Hoffmann - La Roche Inc
340 Kingsland St
Nutley, NJ 07110-1199
(800) 526-6367

For More Information

Contact your doctor or an AIDSinfo Health Information Specialist:

- Via Phone: 1-800-448-0440 Monday - Friday, 12:00 p.m. (Noon) - 5:00 p.m. ET

Isoniazid



For More Information (cont.)

- Via Live Help: http://aidsinfo.nih.gov/live_help
Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET